

in Lateral Position

Critical Incidents

Name:	C Kirk	Observa	tion at start	CRT:	3s
D.O.B.	13/06 (86Y)	RR	12	Temp:	36.2
Address:	(Insert local address)	ETCO2:	1.2	BM:	5.8
		Sats:	95%	Weight:	65
Hospital ID:	1365466214	Heart Rat		Allergy	Penicillin – rash
Ward:	Orthopaedics Background to scenario	BP:	98/56	Specific set up	
An elderly patient is having a hip hemiarthroplasty in the lateral position. After cement application they have a cardiac arrest.			Mannequin – intubated & ventilated Positioned laterally, on theatre trolley IV access, connected to fluid Drugs: vasopressors, used induction drugs Anaesthetic chart		
Required embedded faculty/actors DDP Surgeon and scrub nurse Anaesthetic senior/consultant (handover and leave) Past Medica			Required participants Anaesthetist ODP/Scrub/surgeon can also be participants in MDT sim		
ad a mechani ves in ground f nopping	vears ago, treated medico cal fall (tripped over carpe floor flat, able to potter arc ntures, MP II, normal jaw me	ally, hyperchole et) at home, no ound flat. Has a	sterolaemia, T2D <i>t</i> other injuries. cleaner and relo	tive that helps wi	_
Drugs Home				Drugs Hospital	
amipril 2.5mg (OD			er local protocol)	
andover clinic	DD g OD g OD asked to take over a case al information as above, ir		Antibiotics (as per Femoral block – Occasional vaso icipants ant needs to atte	er local protocol) 20ml 0.25% levob pressors up to no note the processors of the processor of the	oupivacaine ow
amipril 2.5mg (isoprolol 1.25m mvastatin 40m ou have been andover clinic	DD g OD g OD asked to take over a case al information as above, ir	as the consult	Antibiotics (as possible Femoral block – Occasional vaso icipants ant needs to attempt the femoral beautifully, femoral beautifully, femoral beautifully.	er local protocol) 20ml 0.25% levob pressors up to no note the processors of the processor of the	oupivacaine ow
amipril 2.5mg (isoprolol 1.25m imvastatin 40m ou have been andover clinic nd surgery ong	DD g OD g OD asked to take over a case al information as above, in going	e as the consultatubated unever Scenario D Stage 1, Card	Antibiotics (as possible Femoral block – Occasional vascicipants ant needs to attentifully, femoral basinection	er local protocol) 20ml 0.25% levob opressors up to no nd a meeting lock inserted, pos	oupivacaine ow
amipril 2.5mg (isoprolol 1.25m mvastatin 40m ou have been andover clinic nd surgery ong	DD g OD g OD asked to take over a case al information as above, ir	e as the consultatubated unever Scenario D Stage 1, Card	Antibiotics (as possible Femoral block – Occasional vascicipants ant needs to attentifully, femoral basinection	er local protocol) 20ml 0.25% levob opressors up to no nd a meeting lock inserted, pos	oupivacaine ow
amipril 2.5mg (soprolol 1.25mg (soprolol 1.25mg vastatin 40mg vastatin 4	OD g OD g OD asked to take over a case al information as above, in going ventilated. ETCO2 \$1.2. So	s as the consultatubated unever Scenario D Stage 1, Card ats trace – redu	Antibiotics (as por Femoral block – Occasional vasor icipants ant needs to attempt femoral briection liac Arrest ced perfusion \(\) In the control in the ced perfusion \(\) In the c	er local protocol) 20ml 0.25% levob opressors up to no nd a meeting lock inserted, pos	oupivacaine ow sitioned in theatre
amipril 2.5mg (isoprolol 1.25m mvastatin 40m ou have been andover clinic nd surgery ong B Intubated, HR – gradu	OD g OD g OD asked to take over a case al information as above, in going ventilated. ETCO2 \$1.2. So yal bradycardia to 35, ther	scenario D Stage 1, Carc ats trace – redu	Antibiotics (as por Femoral block — Occasional vascicipants ant needs to attendentfully, femoral business and Arrest ced perfusion \$\\$\) Icac Arrest ced perfusion \$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\	er local protocol) 20ml 0.25% levob opressors up to no nd a meeting lock inserted, pos	oupivacaine ow sitioned in theatre
amipril 2.5mg (isoprolol 1.25mg) isoprolol 1.25mg imvastatin 40mg ou have been andover clinic and surgery ong B Intubated, HR – gradu Surgeons of	asked to take over a case al information as above, in going ventilated. ETCO2 \$\frac{1}{2}\$. So all bradycardia to 35, thered closed, Anaesthetised with continue to operate (ceme	sas the consultatubated uneversity Scenario D Stage 1, Carc ats trace – redu PEA. BP 150/2 ith vapour/TIVA ent inserted a fe	Antibiotics (as presented by Femoral block – Occasional vascicipants and needs to attempt for the central by Femoral by F	er local protocol) 20ml 0.25% levob opressors up to no nd a meeting lock inserted, pos	oupivacaine ow sitioned in theatre
amipril 2.5mg (isoprolol 1.25mg) isoprolol 1.25mg imvastatin 40mg ou have been andover clinic and surgery ong IB Intubated, IF Eyes taped Surgeons of Surgeons of Call for he Stop anae Reposition Delegation Treat card Discussion collateral	asked to take over a case al information as above, in going ventilated. ETCO2 \$\frac{1}{2}\$. So yellow a continue to operate (cemerate in condition and the continue to operate in condition and the condition and the continue to operate compression of chance of survival and the continue to operate compression of chance of survival and the condition a	Scenario D Stage 1, Carc ats trace – redu n PEA. BP 50/2 ith vapour/TIVA ent inserted a fe , declare when sions, safe defib	Antibiotics (as por Femoral block – Occasional vascicipants ant needs to attendentfully, femoral business and perfusion decomposition of the composition of the compo	er local protocol) 20ml 0.25% levob pressors up to no nd a meeting lock inserted, pos est essment of cause	oupivacaine ow sitioned in theatre ut, No ROSC
amipril 2.5mg (isoprolol 1.25mg (isoprolol 1.25m	asked to take over a case al information as above, in going ventilated. ETCO2 \$\frac{1}{2}\$. So yeal bradycardia to 35, thered closed, Anaesthetised with a continue to operate (cemerate deterioration in condition lip sthesia, FiO2 1.0 supine of roles – chest compressiac arrest – ALS Principles of chance of survival and	Scenario D Stage 1, Carc ats trace – redu n PEA. BP 50/2 ith vapour/TIVA ent inserted a fe , declare when sions, safe defib	Antibiotics (as por Femoral block – Occasional vascicipants ant needs to attendentfully, femoral businection liac Arrest ced perfusion \$\psi\$ then lost, PEA/company minutes ago) in cardiac arrest perfusion, team assert treatment (No Entire treatment) in the second perfusion of the second perfu	er local protocol) 20ml 0.25% levob pressors up to no nd a meeting lock inserted, pos est essment of cause	oupivacaine ow sitioned in theatre ut, No ROSC
amipril 2.5mg (isoprolol 1.25mg) isoprolol 1.25mg imvastatin 40mg ou have been andover clinic and surgery ong important surgery ong important surgery ong important surgeons of the stop and important card piscussion collateral in the scena	asked to take over a case al information as above, in going ventilated. ETCO2 \$\frac{1}{2}\$. So yellow a continue to operate (cemerate in condition and the continue to operate in condition and the condition and the continue to operate compression of chance of survival and the continue to operate compression of chance of survival and the condition a	scenario D Stage 1, Carc ats trace – redu PEA. BP 50/2 ith vapour/TIVA ent inserted a fe , declare when sions, safe defib planning further nue to a debrie Stage 2, I	Antibiotics (as por Femoral block – Occasional vascicipants and needs to attended to a second perfusion & lac Arrest ced perfusio	er local protocol) 20ml 0.25% levob pressors up to no nd a meeting lock inserted, pos est essment of cause	oupivacaine ow sitioned in theatre ut, No ROSC

Guidelines					
AoA QRH Handbook https://anaesthetists.org/Portals/0/04-12-124225-493	PDFs/QRH/QRH complete April 2022.pdf?ver=2022-				
ANZCA Critical incident debriefing toolkit - https://libguides.anzca.edu.au/criticalincident					
Guidance for Patient Role: N/A					
Opening lines/questions/cues/key responses	Relevant HPC / PMH				
Concerns	Actions				
Guidance for ODP role	Guidance for Scrub team/surgical team				
Opening lines/questions/cues/responses (After incident) Why did this happen? He was doing fine throughout the operation	Initially task focussed, engrossed in surgery until told patient is in cardiac arrest Support with resuscitation efforts				
What did we do wrong He was just like my grandfather/mother, we were talking just before he went to sleep	Support end of life decision making				
Concerns New starter – first cardiac arrest they have experienced					
Competent during the arrest but obviously upset after					
Guidance for Role e.g. ITU/Anaesthetic Senior	Additional challenges				
Expectations/actions Support appropriate for seniority of candidate	Seniority of ODP/other staff Requirement of debrief after incident				
Session Objectives					
Clinical Management of critical incident in non-supine position Management of intra-operative cardiac arrest					

Coordinating team activities, exchanging information, supporting team Planning, following guidance and standards, utilizing available resources

Information gathering, anticipating future steps

Identifying options, MDT decision making

Tell us how you found this simulation scenario resource.

Give us feedback (5 mins) here: https://forms.office.com/e/etz7yZf0aa Or scan the QR code below:



Non-technical skills

Task management Situational awareness

Decision making

Teamworking